



PRELIMINARY GUIDE TO THE IDENTIFICATION OF THE EARLY LIFE STAGES  
OF ARGENTINOIDEI FISH FAMILIES ARGENTINIDAE, MICROSTOMATIDAE,  
AND OPISTHOPROCTIDAE OF THE WESTERN CENTRAL NORTH ATLANTIC




BY

James G. Ditty

U.S. DEPARTMENT OF COMMERCE  
National Oceanic and Atmospheric Administration  
National Marine Fisheries Service  
Southeast Fisheries Science Center  
Galveston Laboratory  
4700 Avenue U  
Galveston, TX 77551

December 2003



**PRELIMINARY GUIDE TO THE IDENTIFICATION OF THE EARLY LIFE STAGES  
OF ARGENTINOIDEI FISH FAMILIES ARGENTINIDAE, MICROSTOMATIDAE,  
AND OPISTHOPROCTIDAE OF THE WESTERN CENTRAL NORTH ATLANTIC**

**BY**

**James G. Ditty**

**U.S. DEPARTMENT OF COMMERCE  
Donald L. Evans, Secretary**

**National Oceanic and Atmospheric Administration  
Conrad C. Lautenbacher, Jr., Under Secretary for Oceans and Atmosphere**

**National Marine Fisheries Service  
William T. Hogarth, Assistant Administrator for Fisheries**

**December 2003**

**This Technical Memorandum series is used for documentation and timely communication of preliminary results, interim reports, or similar special-purpose information. Although the memoranda are not subject to complete formal review, editorial control, or detailed editing, they are expected to reflect sound professional work.**

## NOTICE

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or material mentioned in this publication. No reference shall be made to NMFS or to this publication furnished by NMFS, in any advertising or sales promotion which would imply that NMFS approves, recommends, or endorses any proprietary product or proprietary material mentioned herein or which has as its purpose any intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

This report should be cited as follows:

Ditty, J. G. 2003. Preliminary guide to the identification of the early life history stages of argentinoid fish families Argentinidae, Microstomatidae, and Opisthoproctidae of the western central North Atlantic. NOAA Technical Memorandum NMFS-SEFSC-511, 17 p.

W. J. Richards, Editor. NOAA Fisheries, 75 Virginia Beach Drive, Miami, FL

This report will be posted on the Bethune Cookman College NOAA Cooperative web site later in 2003 at <[www4.cookman.edu/noaa/](http://www4.cookman.edu/noaa/)> and also appear on the SEFSC web site at <[www.sefsc.noaa.gov/](http://www.sefsc.noaa.gov/)>

It will be a chapters entitled Argentinoidei, Argentinidae, Microstomidae, and Opisthoproctidae in "The early life history stages of fishes of the western central North Atlantic".

Copies may be obtained by writing:

the author

NOAA Fisheries

Galveston Laboratory

4700 Avenue U

Galveston, TX 77551

E-mail: <[jim.ditty@noaa.gov](mailto:jim.ditty@noaa.gov)>

or

National Technical Information Center

5825 Port Royal Road

Springfield, VA 22161

(800) 553-6847 or (703) 605-6000

<<http://www.ntis.gov/numbers.htm>>

As currently understood, the sub-order Argentinoidei generally includes four families, the Argentinidae, Bathylagidae, Microstomatidae, and Opisthoproctidae (Begle 1992). Most argentinoids are meso- to bathypelagic in oceanic waters over the outer continental shelf and slope. In members of the Argentinoidei, the dorsal and anal fins develop in the median finfold (rather than along the body wall as is typical of most teleosts) and are connected to the trunk by a series of hyaline strands (Ahlstrom et al. 1984). The pectoral fins develop first, followed by the caudal, dorsal and anal, and pelvic fins. All argentinoids have a deflected terminal gut that hangs free from the body. All regional members of the Argentinoidei have 15 or fewer dorsal rays, and most have fewer than 15 anal rays and an adipose fin. The Argentinoidei have distinctive early life stages with eggs and larvae known for most argentinids, some microstomatids and bathylagids, but few

opisthoproctids. Larvae of the Argentinidae and Bathylagidae are slender, those of Microstomatidae are deeper-bodied, and Opisthoproctidae have both slender and deep-bodied forms. Transformation from the larva to demersal juvenile is protracted in most argentinoids, with pelagic juveniles often retaining their larval pigmentation pattern. Morphological changes generally include a deepening of the body, prolongation of the snout, and eye enlargement. Scale development marks the end of the pelagic juvenile phase. The distinctive ELH stages within each family of the suborder Argentinoidei are unlikely to be confused with those of other taxa. Some taxa have confusing names and a few are listed in Table Argentinoidei 1, otherwise species are discussed in family sections. A list of the species (excluding the Bathylagidae that will be treated later) and their meristics are given in Table Argentinoidei 2.

Table Argentinoidei 1. Synonyms for members of the osmeriform suborder Argentinoidei from the western central North Atlantic, data from Eschmeyer 1998

Family	Common name	Valid name and spelling	Synonyms
Argentinidae	Atlantic argentine	<i>Argentina silus</i>	<i>Salmo silus</i>
		<i>Microstoma microstoma</i>	<i>Microstoma rotundata</i>
Opisthoproctidae	Brownsnout spookfish	<i>Dolichopteryx brachyrhynchus</i>	<i>Bathyllychnops brachyrhynchus</i>
		<i>Dolichopteryx longipes</i>	<i>Aulostoma longipes</i>
		<i>Rhynchohyalus natalensis</i>	<i>Hyalorhynchus natalensis</i>

Table Argentinoidei 2. Meristic data for fishes in the suborder Argentinoidei (except Bathylagidae) from the western North Atlantic. 'NA' indicates no information available.

Species	Vertebrae	Dorsal rays	Anal rays	Pectoral rays	Pelvic rays	Caudal rays (primary+procurent)	Gill rakers (upper-lower or lower only) (+1 at angle in some taxa)	BR	Pyloric caecae	Source
<b>Argentinidae</b>										
<i>Argentina</i>										
<i>brucei</i>	45(44-46)	11(10-12)	(11)12-13	18-19(20)	13-14	19 primary	+7(6)	5	8-10	Cohen & Atsaides 1969
<i>georgei</i>	48(47-50)	11(10-12)	10-11(10-13)	17-18(16-20)	13(12-14)	19 primary	+6(7)	5	8-9	Cohen & Atsaides 1969
<i>silus</i>	64-67	12-13(11)	15-16(13-17)	16-18(15)	12-13	19 primary	6(5-7)+1+12-14(11-15)	6		Schmidt 1906; Moser 1996a Cohen & Atsaides 1969
<i>stewarti</i>	52-53	11(10-12)	12-13	20(18-21)	13-14(15)	19 primary	+6(7)	5	7-9	Cohen & Atsaides 1969
<i>striata</i>	49(47-51)	11-12(10)	12-13(11)	19(18-21)	14(12-15)	10+19+9=38	3(3-4)+6(7)	5	9-14	Cohen & Atsaides 1969
<i>Glossanodon</i>										
<i>polli</i>	?	12-14	11-12(13)	21-22(19-22)	11-12(13)	19 primary	10+1+20-23	5		Cohen 1958
<i>pygmaeus</i>	43-44	11(10-12)	12(11-13)	13(12-14)	10(10-12)	19 primary	10(9-11)+1+22(21-23)	5	8-9	Cohen 1958
<b>Microstomatidae</b>										
<i>Microstoma</i>										
<i>microstoma</i>	44-47	11-13	7-9	7-8	10(9-11)	11+19+11=41	6+1+14	3-4	7	Schmidt 1918; Sanzo 1931a Cohen 1958a; Cohen 1964
<i>Nansenia</i> *										
<i>longicauda</i>	47-50	10-11	10-11	13-14(11-14)	10-11(9)	19 primary	7-8+1+15-18=23-27	4		Moser & Butler 1996
<i>megaloopa</i>	38-39	9-10	8-9	9-10(11)	10-11	19 primary	10-13+1+24-27=36-41	4	7-8	Moser & Butler 1996 Bertelsen 1958
<i>pelagica</i>										
<i>Xenophthalmichthys</i>										
<i>danae</i>	ca. 50	10-12	8-9(10)	7	8-9	10+19+9=38		3	6	Ahlstrom et al. 1984
<b>Opisthoproctidae</b>										
<i>Dolichopteryx</i>										
<i>binocularis</i>	~57-58	13-15	11	13-14	7-9	13+19+13=45		2		Beebe 1932; Beebe 1933
<i>brachyrhynchus</i>	58	10	11	13	8	19 primary		2		Moser 1996d
<i>longipes</i>	41-44	11(10-11)	8-9(10)	11-13	8-9	9-14+19+9-11=40-45		2	4	Beebe 1933; Ahlstrom et al. 1984
<b>Opisthoproctus</b>										
<i>grimaldii</i>	?	12-14	8	11	10	19 primary	20	2		Ahlstrom et al. 1984
<i>soleatus</i>	31	11(10-12)	13-14	9-11	9(10)	19 primary	12	2		Ahlstrom et al. 1984
<i>Rhynchohyalus</i>										
<i>natalensis</i>	?	10-12	8-10	18-20	11-12	19 primary	6+17	4		Masuda et al. 1984; Moser 1996d
<i>Winteria</i>										
<i>telescopa</i>	33-36	8-9	7-9	12-14	7-10			3		Masuda et al. 1984

\* *Nansenia* not likely in our area (Kawaguchi & Butler 1984)

## **ARGENTINIDAE: Argentines or herring smelts**

**By J. G. Ditty**

Adults of the family Argentinidae are silvery and laterally-compressed, with large eyes, a small mouth, toothless jaws, and strong, recurved teeth on the tongue. The pelvic fins are located near mid-body, the anal fin is located far posteriad below the adipose fin, and the caudal fin is forked. Adults are usually <22-cm SL (Moser 1996c). Argentinids have a straight gut that extends to about mid-body and is lined with transverse rugae for most of its length. The head is relatively small with a rounded, blunted anterior profile. Larval pigmentation usually consists of a series of 6-8 bands or patches of pigment ventro-laterally

along the trunk and ends with pigment on the tail. There are five species of *Argentina* and one species of *Glossanodon* in our area (Clarke 2002), but ELH stages are only known for eggs & larvae of *A. silus*, juvenile stages of *A. striata* and *G. polli*. A species account is provided for *A. silus* and an illustration of *G. polli* juvenile is shown in Figure Argentinidae 1. Meristic and distribution characters are given for the family in Table Argentinoidei 2 & Table Argentinidae 1, respectively. Species accounts and illustrations are provided for *Argentina silus*, *A. striata*, and *Glossanodon polli*.

Table Argentinidae 1. Geographic distribution and habitat of the species of the Family Argentinidae

Species	Distribution	Habitat
<b><i>Argentina</i></b>		
<i>brucei</i>	Caribbean & Panama south	Meso- to bathypelagic usually 150-1500 m over mud bottoms
<i>georgei</i>	Atlantic coast of Florida, Dry Tortugas, Caribbean Sea	Meso- to bathypelagic usually 250-450 m over mud & shell bottoms
<i>silus</i>	Northern-most part of the study area	Meso- to bathypelagic usually > 150 m
<i>stewarti</i>	Jamaica & Nicaragua south through southern Caribbean	Meso- to bathypelagic usually 350-550 m
<i>striata</i>		
<b><i>Glossanodon</i></b>		
<i>polli</i>		
<i>pygmaeus</i>		



**MERISTICS**

Vertebrae	
Precaudal	
Caudal	
Total	64-67
Fin rays	
Dorsal	12-13(11)
Anal	15-16(13-17)
Pectoral	16-18(15)
Pelvic	12-13
Caudal	
Principal	19
Adipose Fin	Yes
Gill rakers on first arch	
Upper	6(5-7)+1 angle
Lower	12-14(11-15)
Total	usually 19-21
Branchiostegals	6

**LIFE HISTORY**

Range: northern-most part of study area  
 Habitat: meso- to bathypelagic in oceanic waters usually below 150-m deep  
 ELH pattern: oviparous; pelagic larvae  
 Spawning Season: spring

**LITERATURE**

Ahlstrom et al. 1984, Cohen 1964,  
 Cohen & Atsides 1969, Moser 1996c,  
 Schmidt 1906.

**EARLY LIFE HISTORY DESCRIPTION**

**EGGS:**


Diameter: 3.0-3.5 mm  
 No. of oil globules: one  
 Oil globule diameter: 0.95-1.16 mm  
 Yolk: segmented  
 Shell: smooth  
 Hatch size: 6-9 mm

**LARVAE:**


Length at flexion: ~28-35 mm  
 Length at transformation: >50-mm  
 Sequence of fin development: P<sub>1</sub>, C, D & A, P<sub>2</sub>  
 Pigment: by 17-mm, a characteristic row of 5-6 pigments above gut, & 1 ventrally along caudal peduncle; pigment above & below notochord tip  
 Diagnostic: long, slender body; voluminous finfold; series of pigments above ventrolaterally above hindgut; from other argentinids by the high vertebral and median fin ray counts; 6 branchiostegal rays

**ILLUSTRATIONS**


A). 16.3 mm, B) 19.0 mm, C) 28.0 mm, D) 39.0 mm, E) 50.0 mm SL all redrawn from Schmidt 1906 (see Fahay 1983).




A




B



C



D



E

## ARGENTINIDAE

## *Argentina striata* (Goode & Bean 1896)

### MERISTICS

Vertebrae	
Precaudal	
Caudal	
Total	49(47-51)
Fin rays	
Dorsal	11-12(10)
Anal	12-13(11)
Pectoral	19(18-21)
Pelvic	14(12-15)
Caudal	
Dorsal Secondary	10
Principal	19
Ventral Secondary	9
Total	38
Adipose Fin	Yes
Gill rakers on first arch	
Upper	3(3-4)
Lower	6(7)
Total	9-10
Branchiostegals	5

### LIFE HISTORY

Range: throughout area .

Habitat: meso- to bathypelagic in oceanic waters usually 150-450 m deep over mud bottoms.

ELH pattern: oviparous; pelagic larvae

Spawning: unknown.

### LITERATURE

Ahlstrom et al. 1984, Cohen 1964, Cohen & Atsaiades 1969.

### EARLY LIFE HISTORY DESCRIPTION

EGGS: undescribed.

LARVAE: undescribed


Sequence of fin development: P<sub>1</sub>, C, D & A, P<sub>2</sub>

Pigment: In juveniles, pelvics heavily-pigmented; 11 transverse stripes along body, one on opercle and another across caudal fin; last 7 stripes continuous around ventral side of body but not across dorsum.

### ILLUSTRATIONS

a) Juvenile ca. 20 mm SL, MCZ 66011;

B) Juvenile ca 37 mm SL, MCZ 41865,



A



B

**MERISTICS**

---

Vertebrae	
Precaudal	
Caudal	
Total	Unknown
Fin rays	
Dorsal	12-14
Anal	11-12(13)
Pectoral	21-22(19-22)
Pelvic	11-12(13)
Caudal	
Principal	19
Total	
Adipose Fin	Yes
Gill rakers on first arch	
Upper	10+1 angle
Lower	20-23
Total	31-34
Branchiostegals	5

**LIFE HISTORY**

---

Range: southern-most part of survey area.

Habitat: meso- to bathypelagic in oceanic waters 275-400 m deep.

ELH pattern: oviparous; pelagic larvae.

Spawning: Unknown

**LITERATURE**

---

Ahlstrom et al. 1984, Cohen 1958,  
Poll 1953.

**EARLY LIFE HISTORY DESCRIPTION**

---

EGGS: undescribed.

LARVAE: undescribed


Sequence of fin development: P<sub>1</sub>, C, D & A, P<sub>2</sub>

Pigment: In late stage larvae and juveniles, 8-9 bands laterally along body  
Diagnostic: from most argentinids by the number of D & P<sub>1</sub> fin rays.

**ILLUSTRATIONS**

---


A) 83 mm juvenile from Poll 1953.




Adult microstomatids are generally slender and silvery with a short blunt snout, and dorsal, anal, and pelvic fins placed at or behind mid-body. Most microstomatids are <20-cm SL (Moser & Butler 1996). In *Nansenia*, pelvic fins are located near the termination of the dorsal fin, whereas *Microstoma* and *Xenophthalmichthys* have pelvic fins placed in front of the dorsal fin origin. Microstomatid larvae have a gut that extends to about 75% of body length and has an elongate S-shaped fold in the gut

that lies flat on the left side of the body (Ahlstrom et al. 1984). In microstomatids, the head is bent slightly downward from the longitudinal axis. Most microstomatid larvae have a series of dark pigments above the gut that may extend onto the head and tail. Meristic characters are given in Table Argentinoidei 2. Illustrations of *Microstomus microstomus* eggs, larvae and juveniles are given in Figure Microstomatidae 1.


Figure Microstomidae 1. *Microstomus microstomus*. A) Egg, B) 1.6-mm; C) 6.2-mm; D) 7.2-mm; D) 12.6-mm; E) 15.6-mm; F) 16.8-mm SL, G) *Xenophthalmichthys danae*, 16.5 mm. 1 A-F redrawn after Sanzo (1931); G from Bertelsen 1958.




A




B




C



D



E



F




G



Opisthoproctids are highly specialized, relatively rare fishes from oceanic mid-waters that have an elongate or tubular snout, small terminal mouth, and enlarged pectoral or pelvic fins. The pelvic fins are placed near mid-body and most taxa have tubular, dorsally-directed eyes. Opisthoproctids have a general tendency for delayed metamorphosis and paedomorphosis (Moser 1996d). Larvae of slender-bodied opisthoproctids have an elongate gut while deeper-bodied taxa have a relatively shorter gut. All opisthoproctids

have a sac-like stomach that exits through a constricted pylorus to the intestine (Ahlstrom et al. 1984). Deeper-bodied opisthoproctids have a relatively larger head with a pronounced hump or bend at the nape (Ahlstrom et al. 1984). Opisthoproctids are easily recognized by their well-developed tubular eyes, and precocious pelvic fins (Moser 1996d), and distinctive pigmentation patterns unique to each genus (Ahlstrom et al. 1984). Meristic characters are given in Table Argentinoidei 2.

Figure Opisthoproctidae 1. Illustrations of A) *Opisthoproctus grimaldi*, 14.0 mm; B) *Dolichopteryx binocularis*, 58.0 mm; C) *Dolichopteryx longipes*, 13.4 mm; D) *D. longipes* 27.8 mm, lateral view & E) ventral view of same; G) *Rhynchohyalus natalensis* 23.0 mm. A from Schmidt (1918) as redrawn from Ahlstrom et al. (1984); B from Roule & Angel (1930); C-E from Moser (1996d); G from Bertelsen et al. 1965.



## Argentinoidei Literature

- Ahlstrom, E. H., H. G. Moser & D. M. Cohen. 1984. Argentinoidei: development and relationships. Pages 155-169 in *Ontogeny and Systematics of Fishes*. H. G. Moser et al (eds.). Amer. Soc. Ichthyol. Herpetol. Spec. Publ. (1): 760 p.
- Beebe, W. 1932. Nineteen new species and four post-larval deep-sea fish. *Zoologica* 13(4): 47-107.
- Beebe, W. 1933c. Deep-sea fishes of the Bermuda oceanographic expeditions: Family Argentinidae. *Zoologica* 16(3): 95-147.
- Begle, D. P. 1992. Monophyly and relationships of the argentinoid fishes. *Copeia* 1992: 350-366.
- Bertelsen, E. 1958. The argentinoid fish *Xenophthalmichthys danae*. Dana Rep. (45): 1-8.
- Bertelsen, E., B. Theisen & O. Munk. 1965. On a post-larval specimen, anal light organ, and tubular eyes of the argentinoid fish *Rhynchohyalus natalensis* (Gilchrist and von Bonde). *Vidensk. Meddr. Dansk. Naturh. Foren.* 128: 357-371.
- Cohen, D. M. 1958. A revision of the fishes of the subfamily Argentininae. *Bull. Fla. St. Mus. Biol. Sci.* 3(3): 93-172.
- Cohen, D. M. 1964. Suborder Argentinoidea. Pages 1-70 in *Fishes of the Western North Atlantic*. Sears Fnd. Mar. Res. Mem. 1(Pt 4): 599 p.
- Cohen, D. M. & S. P. Atsides. 1969. Additions to a revision of argentine fishes. *Fish. Bull. U. S.* 68(1): 13-36.
- Eschmeyer, W. N. (ed.). 1998. Catalog of fishes. Center for Biodiversity Research and Information. California Academy of Sciences. Spec. Publ. (1). 3 vols, 2,905 p.
- Fahay, M. P. 1983. Guide to the early stages of marine fishes occurring in the western North Atlantic Ocean, Cape Hatteras to southern Scotian Shelf. *J. Northw. Atl. Fish. Sci.* 4: 1-423.
- Heemstra, P. C. 1986. Family No. 63: Opisthoproctidae. Pages 216-218 in *Smiths' sea fishes*. M. M. Smith & P. C. Heemstra (eds.). Macmillan South Africa. 1047 p.
- Kawaguchi, K. & J. L. Butler. 1984. Fishes of the genus *Nansenia* (Microstomatidae) with descriptions of seven new species. *Contr. Sci.* 352: 22 p.
- Moore, J. A., K. E. Hartel, J. E. Craddock & J. K. Galbraith. 2003. An annotated list of deepwater fishes from off the New England region, with new area records. *N.E. Naturalist* 10(2): 159-248.
- Moser, H. G. 1996c. Argentinidae: Argentines. Page 185-187 in *The Early Stages of Fishes in the California Current Region*. H. G. Moser (ed.) *CalCOFI Atlas* (33): 1505 p.
- Moser, H. G. 1996d. Opisthoproctidae: Spookfishes. Pages 216-223 in *The Early Stages of Fishes in the California Current Region*. H. G. Moser (ed.) *CalCOFI Atlas* (33): 1505 p.

- Moser, H. G. & J. L. Butler. 1996. Microstomatidae: Argentines and pencilfishes. Pages 208-215 in *The Early Stages of Fishes in the California Current Region*. H. G. Moser (ed.). CalCOFI Atlas (33): 1505 p
- Nishimura, S. 1964. Early life history of the deep-sea smelt, *Glossanodon semifasciatus* (Kishinouye) (Teleostei: Clupeida). Part 1. Publ. Seto Mar. Biol. Lab. 13: 349-360.
- Poll, M. 1953. Poissons. III. Téléostéens Malacoptérygiens. Résult. Scient. Expéd. Belge Eaux Côt. Afr. Atlant. Sud. 4(2): 258 p., Plates 1-8.
- Roule, L. & F. Angel. 1930. Larves et alevins de poissons provenant des croisières du Prince Albert I de Monaco. Résultats Camp. Scient. Prince Albert I, 79: 1-148.
- Sanzo, L. 1931. Sottordine: Stomiatoidei. Pages 42-92 + Pls. 4-7 in *Uova, larve e stadi giovanili di Teleostei*. Fauna e Flora del Golfo di Napoli. Stazione Zoologica di Napoli. Monog. 38 (1): 1-177. [In Italian, English translation from Italian in 1969 by Israel Prog. Sci. Transl.; Keter Press, Jerusalem, publisher.].
- Schmidt, J. 1906. On the larval and post-larval development of the argentines *Argentina silus* (Ascan.) and *Argentina sphyraena* (Linne) with some notes on *Mallotus villosus* (O. F. Muller). Meddr. Kommn.. Havunders. Ser. 2(4).
- Schmidt, J. 1918. Argentinidae, Microstomatidae, Opisthoproctidae. Mediterranean Odontostomidae. Rep. Dan. Oceanogr. Exped. Mediterr. 1908-1910 2(A5): 1-40.
- Uyeno, T. 1984. Family Argentinidae. Pages 40-42 in *The Fishes of the Japanese Archipelago*. H. Masuda et al. (eds.), Tokai Univ. Press, Tokyo, Japan. 437 p + Pl. 46.